

An international comparison of patterns of participation in leisure activities for children with and without disabilities in Sweden, Norway and the Netherlands

A. ULLENHAG¹, M.K. BULT^{2,3}, A. NYQUIST⁴, M. KETELAAR^{2,3}, R. JAHNSEN⁵, L. KRUMLINDE-SUNDHOLM¹, L. ALMQVIST⁶, & M. GRANLUND⁷

¹Department of Women's and Children's Health, Karolinska Institutet, Neuropediatric unit, Astrid Lindgren Children's Hospital, Stockholm, Sweden, ²Rudolf Magnus Institute of Neuroscience and Center of Excellence for Rehabilitation Medicine, University Medical Center, Utrecht and Rehabilitation Center De Hoogstraat-Network for Childhood Disability Research in the Netherlands, Utrecht, the Netherlands, ³Network for Childhood Disability Research in the Netherlands, Utrecht, The Netherlands, ⁴Norwegian School of Sport Science, Department of Physical Education and Beitostølen Healthsport Centre, Beitostølen, Norway, ⁵Oslo University Hospital, Department of Neuroscience for children, Rikshospitalet, Oslo, Norway, ⁶CHILD and school of Health, Care and Social Welfare, Mälardalens University, Västerås, Sweden, and ⁷CHILD, School of Health Sciences, Jönköping University, Jönköping, Sweden

(Received 4 May 2012; revised 14 May 2012; accepted 15 May 2012)

Abstract

Purpose: To investigate whether there are differences in participation in leisure activities between children with and without disabilities in Sweden, Norway and the Netherlands and how much personal and environmental factors explain leisure performance.

Methods: In a cross-sectional analytic design, the Children's Assessment of Participation and Enjoyment, CAPE, was performed with 278 children with disabilities and 599 children without disabilities aged 6–17 years. A one-way between-groups ANOVA explored the differences in participation between the countries. Hierarchical multiple regression analysis assessed if age, gender, educational level, living area and country of residence explained the variance in participation. Results: Scandinavian children with disabilities participated in more activities with higher frequency compared to Dutch children. The strongest predictor was country of residence. For children without disabilities, differences existed in informal activities, the strongest predictor was gender.

Conclusion: Differences in school- and support systems between the countries seem to influence patterns of participation, affecting children with disabilities most.

Keywords: Participation, recreation, children, adolescents, disabilities, European comparison

Introduction

Articles 23–30 of the 2006 United Nations (UN) convention on the rights of persons with disabilities declare that children with disabilities should be able to participate on the same terms with others in education, family life, health maintenance, sports and leisure activities [1]. Fauconnier et al. [2] showed, in a cross-sectional European study of children aged 8–12 years with cerebral palsy (CP), that differences in leisure activity patterns of children with CP exist between European countries. One third of the variance could be ascribed to variation between countries. Certain characteristics

of the residential country can be a potential source to differences in participation in leisure activities of children with disabilities. However, differences between countries may also exist for children without disabilities. Comparing children with and without disabilities in different countries provides crucial information on the natural variation between countries and potential differences only present for children with disabilities. Such knowledge may provide insight into how regulations and legislation promote or hinder the accessibility of persons with disabilities to leisure activities.

Participation is defined as a person's involvement in a life situation and can be seen as a result of the

Correspondence: Anna Ullenhag, PT, MSc, PhD student, Health Care Sciences Postgraduate School, Department of Women's and Children's Health, Karolinska Institutet, Neuropediatric unit, Astrid Lindgren Children's Hospital Q2:07, 171 76 Stockholm, Sweden. Tel: +46 (0)73 6620545. Fax: 51777349. E-mail: anna.ullenhag@ki.se

ISSN 1751-8423 print/ISSN 1751-8431 online/12/050369-17 © 2012 Informa UK Ltd.

DOI: 10.3109/17518423.2012.694915

interaction between a person and the environment [3, 4]. Through participation in leisure activities, children are likely to express higher levels of wellbeing, learn new skills and competencies, obtain experience of how society works and have increased opportunities to build friendships [5-7]. Physical leisure activities will help the child to improve functional skills and are linked to health benefits [8-11]. Studies have shown that participation in organized out-of-school leisure activities can benefit a child's academic outcomes, emotions and behaviour [6, 12-14]. Children with disabilities often meet restrictions in their participation due to both personal and environmental factors [5, 15-20]. Pain, mobility problems, communication disorders and intellectual disabilities are examples that can reduce participation [17, 21-23].

For both children with and without disabilities personal factors such as age and gender have an impact on the patterns of participation in leisure activities and when the children enter adolescence the patterns change [24–27]. Participation in skill-based activities decreases, while participation in social activities increases [25–28]. Furthermore, gender differences arise, with girls spending more time in social and skill-based activities than boys. Boys, in contrast, tend to prefer physical activities [5, 12, 27–29].

Environmental factors influencing participation can be related to both the immediate environment of the family and to the wider social environment of the child. The educational level of the parents is related to children's participation in skill-based and physical activities [5, 16, 22, 24, 26, 30, 31]. It is probable that environmental factors have different impacts on the child's pattern of participation in different cultures [2, 32-34]. The way in which children spend their leisure time may vary across countries, influenced as it is by differences in municipal resources, public services and societal and parental expectations [32, 35, 36]. The school structure and school context also influence the child's participation level [6, 37-39]. There is a growing consensus that children with disabilities should be educated within mainstream school settings to emphasize inclusion [4, 30, 36].

Specific environmental factors that have been recognized as obstacles for participation in leisure activities are the social and the physical environments, problems with transportation, fewer municipal resources and public services [2, 15, 22, 23, 35, 36, 40]. Another environmental factor of importance is population density of the child's living area; children and adolescents from rural districts tend to spend more time doing outdoor games, sports and domestic chores, whereas children in urban areas

spend more time in organized sports, in designated play areas or specialized institutions where they are transported by an adult to play and be physically active [29, 41, 42].

The Children's Assessment of Participation and Enjoyment (CAPE) is a self-reported measure of participation assessing the number of leisure activities done and the intensity of the involvement [28, 43]. In this study, the CAPE's diversity and intensity dimensions have been used to systematically study the participation in leisure activities of children with and without disabilities in three European countries.

The objective of this study was to investigate whether participation in leisure activities varies of children with and without disabilities living in Norway, Sweden and the Netherlands. Another objective was to investigate how much of the variation in participation in leisure activities could be explained by environmental variables, after controlling for personal variables.

Methods

Instrument

The CAPE is a self-reported measure consisting of 55 leisure activity items. It is appropriate for children and adolescents with and without disabilities from 6-21 years. The CAPE measures five dimensions of participation: (a) Diversity (the number of activities done in the past 4 months); (b) Intensity (the frequency of participation measured as a function of the number of possible activities within a category) ranging from 1, which corresponds to once in the past 4 months, to 7, which corresponds to once a day or more often; (c) With whom (with whom the child performs the activity most often); (d) Where (where the child performs the activity most often); and (e) Enjoyment (how much the child enjoys doing the activity). These five dimensions of participation each provide three levels of scoring: (i) overall participation score, (ii) scores of participation in 15 formal and 40 informal activities, and (iii) scores reflecting participation divided in five activity types: recreational (12 items), active physical (13 items), social (10 items), skill-based (10 items) and self-improvement activities (10 items). In this study, data distributed by activity types were used concerning diversity and intensity. Several studies have provided indications of reliability and validity of the CAPE outcomes, including studies from Sweden and the Netherlands [28, 43-47]. As the original CAPE questionnaire is in English, forward and backward translations according to international guidelines were carried out in the different countries and

conducted according to the license agreement approved by the publisher.

Sociodemographic variables

Age and gender characteristics were available for all children. Living area was characterized by the number of people living in the residential town. Towns with less than 20 000 residents were characterized as 'rural', towns with more than 20000 residents were characterized as 'urban'. For the Dutch children, the number of residents was determined using postal codes from their home town and a database from Statistics Netherlands and parental educational level was assessed during a phone interview with the parents. For the Norwegian and Swedish children, the number of residents was assessed using the postal code of their residence or the schools and habilitation centres they were recruited from. By using data from Statistics Sweden and Statistics Norway, the corresponding number of residents was determined. The parental educational level was aggregated as 'non-university level' or 'university level' based on the study questionnaire for the Norwegian parents and on the reports of the parent's occupation obtained from a database from Statistics Sweden.

Procedures

A cross-sectional analytic design was applied. Somewhat different data collection methods were used in the different countries. In Sweden, the CAPE was performed with 55 children with disabilities aged 6-16 years old (M=11.7 years)SD = 2.0), recruited from 13 habilitation centres situated in rural and urban areas in the central part of Sweden. All the participants filled in the CAPE questionnaire at home with help from parents when needed (Tables II and III) (Ullenhag et al. submitted). In addition, 337 typically-developing children aged 6-17 years (M=12.0, SD=2.0) responded to the CAPE. The children were recruited from six regions/schools situated in the central part of Sweden, including coastal and snowy landscapes, rural and urban areas and different socioeconomic districts. The children completed the CAPE questionnaire in the classroom with one of the researchers available to answer questions. The youngest children (aged 6-7 years) completed the CAPE questionnaire at home, in case they needed the assistance of a parent [47].

The 165 Norwegian children with disabilities aged 8–17 years (M=11.8, SD=2.5) who participated were recruited during their rehabilitation period at Beitostølen Healthsport Center (BHC). The physiotherapists working at BHC delivered the CAPE to the families. All the participants answered the CAPE

questionnaire together with their families and a physiotherapist was available to answer questions. A total of 107 typically-developing children aged 7–14 years (M=11.1, SD=2.5) also participated. These children were recruited from five schools situated in urban and rural areas of southeastern Norway with different socioeconomic districts. The children completed the CAPE questionnaire in the classroom with one of the researchers and the teacher available to answer questions.

From the Netherlands, 74 children with disabilities participated aged 6-18 years (M=12.0,SD=3.4). The children were recruited from two schools for special education and from a rehabilitation centre. The children were eligible if they were able to complete the CAPE with or without assistance. The CAPE questionnaire was completed in school or at home in a one-on-one session together with a research assistant. A convenience sample of 158 children without disabilities aged 6-18 years (M=11, SD=3.1) was recruited from Dutch regular schools in different regions of the Netherlands. Children younger than 12 years completed the CAPE in a one-on-one session with a research assistant. Children aged 12-18 years filled out the CAPE in the classroom with a research assistant available for assistance if needed [45].

Ethical considerations

The Regional Committee for Medical Research Ethics, in Uppsala, Sweden, approved the study of Swedish participants Reference No. 2008/394. The Norwegian study was approved by the Regional Committee for Medical Research Ethics, South-East Norway (REK-South East-A no: S-08658a) and the Data Inspectorate (no: 20095). The University Medical Center Utrecht, in the Netherlands and the management of all participating local schools approved the Dutch study. For all children, informed consent was given by parents and children.

Data analysis

Participants were grouped by age, gender, country of residence, the mothers' level of education, rural or urban living areas. The analyses were based on the sum of diversity scores for each of the five activity types. To simplify the analysis, the original eight CAPE categories of intensity were merged into three new categories representing low, regular or high intensity of participation in leisure activities; (i) Seldom/never (original intensity scale 0 = never, 1 = once/4 months, 2 = twice/4 months), (ii) Regularly (original intensity scale 3 = once/month, 4 = 2-3 times/month) and, finally, (iii) Often (original intensity scale 5 = once/week, 6 = 2-3 times/week, and 7 = once/day or more often).

Table I. Demography of children with and without disabilities.

· ·		Sweden	ue .	The Netherlands	lands	Norway	ay
	1	Children	Children	Children	Children	Children	Children
		without	with	without	with	without	with disabilities,
		disabilities,	disabilities,	disabilities,	disabilities,	disabilities,	(%) u
		(%) u	(%) u	(%) u	n (%)	(%) u	
Participants		. 337*	55	158	74	104	149
Age	6-8 years	35 (10.4).	14 (25.5)	51 (32.2)	20 (27.0)	8 (7.7)	34 (22.8)
)	9-12 years	189 (56.3)	21 (38.2)	40 (25.3)	19 (25.7)	50 (48.1)	49 (32.9)
,	13-18 years	112 (33.3)	. 20 (36.4)	67 (42.4)	35 (47.3)	46 (44.2)	66 (44.3)
Mean (min-max), SD		12.0 (6–17), 2.0	11.7 (6–16), 2.7	11.0 (6–17), 3.1	12.0 (6–18), 3.4	11.1 (7–14), 2.5	11.9 (8–18), 2.5
Sex	Male	164 (48.7)	29 (52.7)	72 (45.6)	38 (51.4)	47 (45.2)	82 (55.0)
	Female	173 (51.3)	26 (47.3)	86 (54.4)	36 (48.6)	57 (54.8)	67 (45.0)
Rural area ≥ 20000	•	76 (22.6)	15 (27.3)	58 (36.7)	12 (16.2)	50 (48.1)	78 (52.3)
Urban area $\leq 21~000$		261 (77.4)	40 (72.7)	87 (55.1)	59 (79.7)	54 (51.9)	71 (47.7)
Missing				13 (8.2)	3 (4.1)		
Mothers educational level	*	119 (35.3)	19 (34.5)	25 (15.8)	45 (60.8)	60 (57.7)	· 76 (51.0)
less than university			•				
University level		144 (42.7)	20 (36.4)	41 (25.9)	26 (35.1)1	39 (37.5)	71 (47.7)
Missing		74 (22.0)	16 (29.1)	92 (58.2)	3 (4.1)	5 (4.8)	2 (1.3)

Clinical condition/ Sweden, The Netherlands, Norway, Clinical condition diagnosis of children n (%) n (%) n (%) Central Nervous system Cerebral palsy or related 24 (43.6) 28 (37.8) 62 (37.6) **GMFCS GMFCS GMFCS** Ι 5 (20.8) 12 (42.9) 12 (21.8) II 4 (16.7) 78 (28.6) 21 (38.2) Ш 5 (20.8) 6(21.4)15 (27.3) īν 4 (16.7) 1 (3.6) 7 (12.7) 0 (0) 1 (3.6) Spina bifida, spinal cord or related 8 (14.5) 6 (8.1) 16 (10.7) Motor delay 1 (1.8) 4 (5.4) Developmental delay 17 (11.4) 10 (13.5) Aquired brain injury 3 (4.1) 4 (2.7) Other CNS 2(3.6)12 (16.2) 29 (19.5) Musculoskeletal Neuromuscular 5 (9.1) 6 (8.1) 5 (3.4) skeletal 8 (14.5) 5 (6.8) Other musculoskeletal 2(3.6)Arthritis 6 (8.1%) 9 (6.0) Other 4 (7.3) 14 (9.4) Missing 1(1.8)

Table II. Demography of children with and without disabilities.

A Company Andrews on residence of a graph base of a second

4 4 4 3 18 18 18

The overall percentage of activities done seldom/never, regularly and often of five activity types were calculated for each participant as follows; The number of activities done in each category, seldom/never, regular or often was divided by the total number of activities in each activity scale and the ratio was multiplied by 100. For example if one child had done three of 13 recreational activities often, the overall percentage of recreational activities done often was $3/13 \times 100 = 23\%$. The use of percentages instead of absolute numbers enabled comparison of five activity types with different numbers of items.

One-way between-groups analysis of variance tests (ANOVA) were conducted to explore the impact of country of residence on the percentage of activities done: seldom/never, regularly and often and on diversity for each of the five activity types. Post-hoc comparisons using the Tukey test were performed when a significant difference between groups was present. The one-way between-groups ANOVA was also used to analyse if there were significant differences between the three countries regarding age. Chi-squared analyses were calculated to analyse if differences existed between the countries regarding gender, the educational level of mothers and the geographical area of living.

Hierarchical multiple regression analysis tests were used to assess the ability of the three independent environmental variables: educational level, geographical living area and country of residence, to predict the outcome scores of diversity and intensity in five activity patterns of participation, after controlling for the influence of the personal variables age and gender. The rationale for using hierarchical regression

analysis was to evaluate personal and environmental factors separately, as personal factors are less contextdependent. Since the personal factors; age and gender are known to influence the pattern of participation in leisure activities, regardless of nationality it was of interest to evaluate if environmental factors could further explain the variance of participation. Preliminary analyses were performed to ensure no violation of the assumptions of normality, linearity, multi-co-linearity and homoscedasticity. Age and gender were entered in Step 1, while educational level, geographic area and country of residence with dummy variables for each country were entered in Step 2. Statistical analyses were performed using the SPSS for Windows software program (version 19.0) and the chosen alpha level for all analyses was p = 0.05.

Results

Demographic differences

Demographic variables are described in Tables I and II. For children with disabilities, no significant differences were found between the three countries except for differences in rural or urban living areas. A significantly higher proportion of children with disabilities from Sweden (73%) and the Netherlands (80%) were living in urban areas compared to the Norwegian children (48%) with disabilities; $\chi^2(2, n=275), p=0.000$.

For children without disabilities there were significant differences in educational level of the mothers $\chi^2(2, n=429), p=0.007)$ and in the geographical

Table III. Children with disabilities-diversity and intensity participation in leisure activities.

		Sweder	ı		Netherlar	nds		Norway	
	n	M (SD)	Range	n	M (SD)	Range	n	M (SD)	Range
No. of activities done			•						
Recreational activities*(12)	55	7.1 (2.7)		74	6.4 (2.7)		149	7.3 (2.4)	
Physical activities**(13)	55	3.5 (2.2)		74	2.4 (1.5)		149	3.2 (1.8)	
Social activities**(10)	55	6.1 (1.8)		74	4.3 (2.0)		149	6.6 (1.8)	
Skill-based activities**(10)	55	2.7 (1.7)		74	1.5 (1.5)		149	2.2 (1.4)	
Self-improvement activities**(10)	55	4.3 (1.6)		74	3.1 (1.8)		149	4.3 (1.5)	
No. of activities done seldom									
Recreational activities*		39.6%	0-92%		46.9%	0-100%		39.3%	0–83%
Physical activities**		72.5%	38-100%		81.4%	54-100%		75.3%	38-100%
Social activities**		38.4%	0–80%		56.6%	10-100%		40.0%	0–80%
Skill-based activities**		72.4%	40-100%		84.6%	30-100%		78.3%	30-100%
Self-improvement activities**		56.3%	20–90%		71.1%	30–100%		57.1%	20–90%
No. of activities done regularly		•							
Recreational activities**		17.4%	0-42%		8.3%	0–33%		16.2%	0-58%
Physical activities**		11.5%	0-38%		3.0%	0–15%		9.5%	0-46%
Social activities**		29.3%	0-70%		17.9%	0-50%		28.0%	0–70%
Skill-based activities**		8.7%	0-30%		2.3%	0-20%		5.6%	0-30%
Self-improvement activities**		15.7%	0–40%		7.7%	0–50%		12.1%	0–50%
No. of activities done often									
Recreational activities		43.0%	0-100%		44.8%	0–92%		44.5%	8-83%
Physical activities		15.9%	0–38%		15.6%	0-46%		15.2%	0-62%
Social activities**		32.3%	0-60%		25.5%	0–80%		38.1%	0–70%
Skill-based activities		18.9%	0-50%		13.1%	0–70%		16.1%	0-60%
Self-improvement activities**		28.0%	0-50%		21.2%	0–60%		30.9%	0-80%

^{*}p < 0.05; **p < 0.01.

Recreation 12 items; Physical 13 items; Social 10 items; Skill-based 10 items; Self-improvement 10 items.

living area $\chi^2(2, n=586)$, p=0.000. A significantly higher percentage of the mothers in Sweden (43%) and in Norway (38%) had a university education compared to mothers in the Netherlands (26%). There was a significantly higher proportion of the Swedish children (77%) living in urban areas compared to children in Norway (52%) and in the Netherlands (55%).

Differences between countries in diversity outcome

One-way between-groups ANOVA indicated significant differences due to country of residence on the diversity of activities in the five activity types for children with disabilities. *Post-hoc* comparisons showed that the Norwegian and Swedish children with disabilities performed a significantly higher number of activities than the children from the Netherlands in all activity types, except for recreational activities. In recreational activities there was a significant difference between Norway and the Netherlands, with the Norwegian children participating in a higher number of activities (Table III).

For children without disabilities, Swedish and Norwegian children participated in a higher number of social activities than children from the Netherlands. On the other hand, the Dutch children participated in a higher number of recreational activities than children in Sweden and Norway. No significant differences were found in the other activity types (Table IV).

Differences between countries in intensity outcome

Seldom/never. The ANOVAs revealed significant differences in intensity never/seldom of children with disabilities due to country of residence. Children in the Netherlands more commonly performed recreational activities seldom/never (47%) than children with disabilities from Norway (39%), p < 0.05. Children from the Netherlands to a higher degree participated seldom/never in physical, social, skill-based and self-improvement activities compared to Norwegian and Swedish children with disabilities, p < 0.05 (Table III).

For children without disabilities, post-hoc comparison revealed that the children from the Netherlands more commonly participated in social activities seldom/never than Swedish and Norwegian children. In physical activities, children from the Netherlands participated more seldom/never compared to Swedish children. The opposite was found concerning recreational activities, where children in Sweden and Norway participated more seldom/never than

Table IV. Children without disabilities-diversity and intensity participation in leisure activities.

		Sweden			The Netherlar	nds		Norway	
	n	M (SD)	Range	n	M (SD)	Range	n	M (SD)	Range
No. of activities done					-	,			
Recreational activities**(12)	337	5.8 (2.6)	N.	158	7.1 (2.3)		104	5.9 (2.5)	
Physical activities(13)	337	4.0 (2.2)		158	3.5 (1.7)		104	3.9 (2.2)	
Social activities**(10)	337	6.1 (1.8)		158	5.3 (2.0)		104	6.3 (1.9)	
Skill based activities(10)	336	2.0 (1.7)		158	2.1 (1.5)		104	2.5 (2.0)	
Self-improvement activities(10)	337	4.0 (1.8)		158	4.3 (2.0)		104	4.4 (1.7)	
No. of activities done seldom									
Recreational activities**		49.9% (21.7%)	0-100%		41.1% (19.3%)	0-92%		51.2% (20.7%)	8-92%
Physical activities**		68.3% (17.6%)	13-100%		72.9% (13.3%)	42-100%		69.7% (16.8%)	23-100%
Social activities**		37.6% (17.9%)	0-100%		47.1% (20.2%)	10~90%		37.1% (19.4%)	0-100%
Skill based activities		79.0% (18.2%)	0-100%		79.4% (15.1%)	30-100%		74.8% (20.0%)	10-100%
Self-improvement activities		58.8% (18.8%)	0-100%		57.1% (20.0%)	10-100%		56.4% (16.8%)	20100%
No. of activities done regularly									
Recreational activities*		16.5% (13.4%)	0-58%		14.8% (10.8%)	0-42%		12.7% (11.6%)	0-50%
Physical activities**		9.9% (10.4%)	0-62%		5.2% (6.5%)	0-31%		7.2% (7.5%)	0-31%
Social activities*		21.4% (13.8%)	0-70%		24.6% (15.3%)	0-80%		19.6% (15.1%)	0-80%
Skill based activities		4.7% (7.4%)	0-40%		4.6% (6.3%)	0-20%		6.5% (10.7%)	0-70%
Self-improvement activities		11.9% (10.9%)	0-50%		11.5% (9.9%)	0–40%		10.1% (9.4%)	0-40%
No. of activities done often									
Recreational activities**		33.6% (18.7%)	0-92%		44.0% (20.2%)	8-100%		36.1% (18.5%)	0-83%
Physical activities		21.8% (14.3%)	0–75%		22.0% (12.0%)	0-54%		23.2% (15.5%)	0-69%
Social activities**		40.0% (15.5%)	0-90%		28.3% (13.9%)	0-80%		43.1% (18.2%)	0-80%
Skill based activities		16.3% (16.7%)	0-100%		16.1% (14.3%)	0-60%		18.6% (17.5%)	0-90%
Self-improvement activities*		29.3% (15.2%)	0–80%		31.3% (15.7%)	0-80%		33.5 (15.7%)	0-80%

^{*}p < 0.05; ** p < 0.01.

Recreation 12 items; Physical 13 items; Social 10 items; Skill-based 10 items; Self-improvement 10 items.

children in the Netherlands. There were no significant differences in the other activity types due to country of residence (Table IV).

Regularly. Significant differences were found due to country of residence in intensity regular. A post-hoc comparison revealed that Swedish and Norwegian children with disabilities participated in a significantly higher percentage of the recreational, physical. social. skill-based and self-improvement activities regularly than children from Netherlands (Table I).

Among the children without disabilities, Swedish children participated in a higher percentage of recreational activities regularly compared to the Norwegian children and they also participated in physical activities more regularly than children from Norway and the Netherlands. Further, the children from the Netherlands participated in a higher percentage of social activities regularly than the Norwegian children (Table IV). No significant differences were found due to country of residence in the other activity types.

Often. The ANOVA revealed significant differences in intensity 'often' due to country of residence.

Swedish and Norwegian children with disabilities participated in a significantly higher percentage of the social activities often than children with disabilities from the Netherlands. In self-improvement activities, Norwegian children participated in a higher percentage of the activities often compared to children in the Netherlands. No significant differences were found in recreational, physical and skill-based activities between the children due to country of residence (Table III).

Children without disabilities from the Netherlands performed a higher percentage of the recreational activities often compared to children from Sweden and Norway. However, in social activities the pattern was reversed. The children from Norway participated in self-improvement activities more often than Swedish children. No significant differences were found in physical and skill-based activities between the children due to country of residence (Table IV).

Personal and environmental variables predicting outcome scores of diversity

The environmental variables hypothesized to have an effect on the diversity scores in the five activity types were: the educational level of mothers, geographic living area and country of residence. The

Table V. The strongest model and predictor of variance in diversity outcome of five activity types for children with and without disability.

		,	Children with disa	ability		Childre	en without	disabilities
Activity type	Step 1	Step 2	Sig. F change	Strongest variable (Correlation part ²)	Step 1	Step 2	Sig.F change	Strongest variable (Correlation part ²)
Recreation	24%	27%	0.066	Age (22.6%)	7%	11%	0.000	Age (5.1%)
Physical	6%	12%	0.000	Gender (4.5%)	7%	9%	0.053	Gender (6.7%)
Social	2%*	25%	0.000	Country NL (18.2%)	3%	8%	0.000	Country NL (4.1%)
Skill-based	7%	15%	0.000	Gender (4.8%)	10%	13%	0.015	Gender (9.8%)
Self-improvement	2%*	12%	0.000	Country NL (7.1%)	12%	12%	0.754	Gender (10%)

Step 1 = Age and gender; Step 2 = mothers educational level, rural/urban living area and country of residence as three dummy; SW, Sweden; NL, the Netherlands; NO, Norway.

personal variables age and gender were added as control variables in the hierarchical regression models. After controlling for the personal variables of children with disabilities, the environmental variables could additionally explain a significant proportion of the variance in all activity types except for recreational activities. The total variance in diversity scores explained by the environmental variables varied between 12.0–26.7%.

For children without disabilities the environmental variables in the hierarchical multiple regression analysis explained less of the variance, 8.3-12.7% of the diversity outcome scores of five activity types. The environmental variables significantly explained the variance in recreational, social and skill-based activities after controlling for age and gender. The main predictor was the same for both children with and without disabilities in all activity types except for in self-improvement activities. In recreational, physical and skill-based activities, personal variables mainly contributed to the diversity. Age was the main predictor in recreational activities and gender in physical and skill-based activities. The environmental variable country of residence was the main predictor in social activities and also in selfimprovement activities for children with disabilities. Gender was the main predictor in self-improvement for children without disabilities. Results in detail are presented in Table V and in the Appendix.

Personal and environmental variables predicting outcome scores of intensity

Overall, the hierarchical multiple regression analysis showed that the personal and environmental independent variables could explain 8.2–26.7% of the variance of intensity (activities done seldom/never, regular or often) in the five activity types for children with disabilities. The environmental variables could best explain the variance in activities done seldom/never and regularly. The country of residence was

the main predictor in nine of 15 (60%) intensity activity types and the strongest predictor in all activities done on a regular basis and in social and self-improvement intensity outcome scores. Age was the strongest predictor in recreational activities and gender was the strongest predictor in physical and skill-based activities. Results in detail are presented in Table VI and in the Appendix.

For children without disabilities the hierarchical multiple regression analysis could predict less of the variance, between 3.2–14.9% of intensity scores in five activity types. Results in detail are presented in Table VI and in the Appendix.

The environmental variables could best explain the variance in activities done seldom/never. Gender was the strongest predictor in nine of 15 (60%) intensity activity types and the main predictor in physical, skill-based and self-improvement activities. Country of residence was the main predictor in recreational and social activities. In summary the same main predictors could explain the greater part of the variance in physical, social and skill-based activities for children with and without disabilities. In general, the environmental variable, country of residence was more frequently the main predictor in intensity outcome scores for children with disabilities; while for children without disabilities the personal variable, gender was most often the main predictor in diversity and intensity outcome scores.

Discussion

The results of this study indicate that there are differences between countries in patterns of participation in leisure activities for children with disabilities both regarding diversity and intensity. For children without disabilities there were mainly differences between the countries in recreational and social activities. The results show that the environmental variable, country of residence, was

^{*}non-significant.

Table VI. Summary of hierarchical regression analysis. The strongest model and predictor of variance in intensity outcome of five activity types for children with and without disabilities explained of variance in percentages.

			Ch	ildren w	ith disabilities		Child	ren v	without d	isabilities
Activity type	Step 1	Step 2	R^2	Sig.F change	Strongest variable (Correlation part ²)	Step 1	Step 2	R^2	Sig.F change	Strongest variable (Correlation part ²)
Recreation										
Seldom/never	24%	27%		0.076	Age (22.6%)	7%	11%		0.003	Age (5.5%)
Regular	4%	15%		0.000	Country NL (8.2%)	1%*	3%		0.049	Country NO (1.4%)
Often	18%	19%		0.744	Age (17.0%)	5%	10%		0.000	Country NL/Age (4.1%/3.9%)
Physical										
Seldom/never	6%	14%		0.000	Gender/living (4.4%/3.3%)	7%	10%		0.022	Gender (6.7%)
Regular	$0.5\%^{1}$	14%		0.000	Country NL (6.2%)	0.5%*	6%		0.000	Country NL (4.8%)
Often	6%	8%		0.172	Gender (4.8%)	8%	9%		0.469	Gender (7.6%)
Social										
Seldom/never	2%	24%		0.000	Country NL (17.6%)	3%	9%		0.000	Country NL (4.9%)
Regular	$0.5\%^{1}$	12%		0.000	Country NL (7.8%)	3%	4%		0.164	Gender (1.6%)
Often	2%1	14%		0.000	Country NL (7.7%)	2%	15%		0.000	Country NL (11.3%)
Skill-based										
Seldom/never	7%	15%		0.000	Gender (4.8%)	9%	11%		0.055	Gender (8.6%)
Regular	0.5%¹	10%		0.000	Country SW/NL (2.9%/2.2%)	2%	4%		0.245	Gender (2.0%)
Often	6%	10%		0.079	Gender (5.3%)	7%	9.0	%.	0.013	Gender (6.6%)
Self-improvement	:				•					
Seldom/never	1%*	15%		0.000	Country NL (10.0%)	12%	12%		0.913	Gender (9.8%)
Regular	0%* ¹	8%		0.000	Country SW/NL (2.0%/1.8%)	2%	3%¹		0.597	Gender (2.0%)
Often	2%*	10%		0.000	Country NL(7.8%)	10%	11%		0.505	Gender (7.8%)

Step l = Age and gender; Step 2 = mothers educational level, rural/urban living area and country of residence as three dummy; SW, Sweden; NL, the Netherlands; NO, Norway.

the strongest predictor of variance in all activity types performed on a regular basis for children with disabilities. This concerned both the diversity and the intensity outcome scores, in particular in social and self-improvement activities. On the contrary, for children without disabilities fewer differences between the countries were found and mainly in recreational physical and social activities.

In general, Scandinavian children with disabilities appear to have a higher level of participation in leisure activities compared to the Dutch children with disabilities. These results are in line with the cross-sectional European study of participation in children aged 8-12 years with CP, where children from Denmark had the highest level of participation [2]. The reason for the differences cannot be determined from this study alone, but the prominent difference between Scandinavian countries and the Netherlands regarding children with physical disabilities may concern the education systems. Studies have shown that children with disabilities educated within mainstream schools have a higher level of participation in some recreational activities compared to children in special schools [2, 4, 30]. There is also evidence to

suggest that to include children with disabilities in regular classes increases their opportunities to gain the same social participation and theoretical knowledge as children without disabilities [30, 37, 48]. In Sweden and Norway, children with disabilities are by law [49, 50] included into mainstream schools and only children with blindness, deafness, mental retardation (IQ level below 70), autism or similar conditions [49, 50] are segregated into special schools. In the Netherlands, 62% of the children with special educational needs attend segregated schools, compared to less than 4% in Norway and Sweden (SNE Country data, 2010) [51], which may be one of the explanations of the differences in the participation level. In addition, the administration of the CAPE requires that the activities are done outside of the school curriculum. This might indicate that the Dutch children with disabilities participate in corresponding activities such as swimming, horseback riding, etc., during the school hours and, since the activities are within the school curriculum they will not be registered as leisure activities in the CAPE. Another factor that may affect the results concerns the cultural applicability of the item content in the CAPE.

^{*}non-significant.

Other explanations of the differences in the participation patterns might be that Sweden and Norway have public after-school recreational centres attended all weekdays by both children with and without disabilities [36, 52]. These centres provide the children with opportunities to be engaged in social and recreational activities. Transportation and accessibility may be obstacles for participation in leisure activities for children with disabilities. Sweden and Norway might have a more generous national policy of transportations for children with disabilities than other European countries, which probably facilitates participation in leisure activities (Social security Services 1997, 2002; Regulation no 290, Available transport) [36, 52]. In Sweden there are also policies for social care and support services and national resources for providing information about assistive technology. Different policies and legislation directed at assistive technology, equality of information, support and welfare services, etc., might be other important factors that influence the participation profile of children in different countries [2, 36].

For children without disabilities only minor differences between countries could be seen. Children from Sweden and Norway participated in unorganized social and recreational activities with a higher intensity and diversity compared to the Dutch children. Again, this may be an effect of different school systems. In general, the Swedish and Norwegian children have more unscheduled out-of-school time compared to west European children and this may partly explain the results.

Gender was the strongest predictor of variance in both diversity and intensity outcome scores for children without disabilities. This result is in accordance with other studies showing that boys perform more physical activities while girls spend more time in social and skill-based activities [5, 12, 27–29]. The pattern of societal, parental and peer expectations influence the participation in leisure activities and needs further consideration. It is of special concern that girls' participation in physical activities tends to decrease with increasing age, since involvement in physical activities is crucial for health and well-being.

A large sample of children with and without disabilities from the three countries was included in the study. Even though the inclusion criteria were comparable in all three countries, a limitation is that one cannot know if the children's functional levels were equivalent. Studies of children with CP have reported that a lower level of functioning is correlated with lower participation level [17, 20, 23, 27]. There are studies indicating that type of disability is of less importance for the pattern of participation than severity of impairment [53, 54]. However,

functional classifications for gross motor function, manual ability and communication such as the GMFCS [55], the Manual Ability Classification System, MACS [56] and the Communication Function Classification System, CFCS [57] are available only for children with CP. Evaluating if these classifications also could be used for other populations of individuals with functional limitations would be valuable. Other factors, such as social skills and autonomy [53, 54], also need further consideration when selecting assessment instruments/classifications and research methodology.

Another potential limitation of this study concerns that data collection was done by different therapists and researchers in each country, which may have introduced systematic differences. This risk was minimized by training the involved therapists in the Netherlands and Norway. In Sweden, the children with disabilities filled in the CAPE at home together with a parent and 53-81% of the families, respectively, found the CAPE intensity and diversity questions rather easy to answer. The Swedish and Norwegian children without disabilities filled in the CAPE in their classrooms with a researcher available to answer questions. However, all the participants responded to the same instrument, the CAPE has reported sufficient interinterviewer reliability and the intra-class correlation coefficient for the intensity scores ranged from 0.66-0.83 [45].

The results from this study revealed that the personal and environmental independent variables in the model could explain only 3–27% of the variance of the diversity and intensity outcome scores. These findings are consistent with similar studies by King et al. [27], Palisano et al. [23] and Imms et al. [38]. The larger unexplained variance in this study indicates that the model requires additional determinants, such as physical accessibility, availability of transportation, community programmes and the family's preferences and orientation to social and cultural leisure activities. The functional level of all the children with disabilities may also be important variables. All these factors are potential determinants that were not included in this model.

In conclusion, this study found that environmental variables considerably influenced the patterns of participation for children with disabilities. As environmental factors often are amenable to state intervention, further analysis of the physical accessibility, community programmes and the family's preferences to social and cultural leisure activities, etc., and the causes of nationality heterogeneity are essential to provide evidence for changes to national legislation and policies that promote participation for children with disabilities.

Acknowledgements

We would like to thank all the children, parents, therapists and research assistants who participated in this project. We acknowledge also the Johanna Kinderfonds and the Stichting Rotterdams Kinderrevalidatie Fonds and Adriaanstichting for founding the Dutch part of this research.

Declaration of interest: The authors report no conflicts of interests. The authors alone are responsible for the content and writing of the paper.

References

- United Nations. Convention on the rights of persons with disabilities. New York: United Nations; 2006.
- Fauconnier J, Dickinson HO, Beckung E, Marcelli M, McManus V, Michelsen SI, Parkes J, Parkinson KN, Thyen U, Arnaud C, et al. Participation in life situations of 8-12 year old children with cerebral palsy: Cross sectional European study. British Medical Journal 2009;338:b1458.
- World Healt Organization WHO. The international classification of functioning, disability and health. Geneva, Switzerland: WHO; 2001.
- Hoogsteen L, Woodgate RL. Can I play? A concept analysis
 of participation in children with disabilities. Physical &
 Occupational Therapy in Pediatrics 2010;30:325-339.
- Law M, King G, King S, Kertoy M, Hurley P, Rosenbaum P, Young N, Hanna S. Patterns of participation in recreational and leisure activities among children with complex physical disabilities. Developmental Medicine and Child Neurology 2006;48:337-342.
- Feldman AF, Matajsko JJ. The role of school-based extracurricular activities in adolescent development: A comprehensive review and future directions. Review of Educational Research 2005;75:159-210.
- Solish A, Perry A, Minnes P. Participation of children with and without disabilities in social, recreational and leisure activities. Journal of Applied Research in Intellectual Disabilities 2010;23:226-236.
- Verschuren O, Ketelaar M, Gorter JW, Helders PJ, Uiterwaal CS, Takken T. Exercise training program in children and adolescents with cerebral palsy: A randomized controlled trial. Archives of Pediatric and Adolescent Medicine 2007;161:1075-1081.
- Janssen I, Leblanc AG. Systematic review of the health benefits of physical activity and fitness in school-aged children and youth. International Journal of Behavioral Nutrition and Physical Activity 2010;7:40.
- Dodge T, Lambert SF. Positive self-beliefs as a mediator of the relationship between adolescents' sports participation and health in young adulthood. Journal of Youth and Adolescence 2009;38:813–825.
- Ploughman M. Exercise is brain food: The effects of physical activity on cognitive function. Developmental Neurorehabilitation 2008;11:236–240.
- Simpkins SD, Ripke M, Huston AC, Eccles JS. Predicting participation and outcomes in out-of-school activities: Similarities and differences across social ecologies. New Directions of Youth Development 2005;105:51-69, 10-11.
- Dahan-Oliel N, Shikako-Thomas K, Majnemer A. Quality of life and leisure participation in children with neurodevelopmental disabilities: A thematic analysis of the literature. Quality of Life Research 2012;21:427-39.

- 14. Chomitz VR, Slining MM, McGowan RJ, Mitchell SE, Dawson GF, Hacker KA. Is there a relationship between physical fitness and academic achievement? Positive results from public school children in the northeastern United States. Journal of School Health 2009;79:30-37.
- Parkes J, McCullough N, Madden A. To what extent do children with cerebral palsy participate in everyday life situations? Health and Social Care in the Community 2010;18:304-315.
- King G, Law M, King S, Rosenbaum P, Kertoy MK, Young NL. A conceptual model of the factors affecting the recreation and leisure participation of children with disabilities. Physical & Occuptional Therapy in Pediatrics 2003;23:63-90.
- Majnemer A, Shevell M, Law M, Birnbaum R, Chilingaryan G, Rosenbaum P, Poulin C. Participation and enjoyment of leisure activities in school-aged children with cerebral palsy. Developmental Medicine and Child Neurology 2008;50:751-758.
- Engel-Yeger B, Jarus T, Anaby D, Law M. Differences in patterns of participation between youths with cerebral palsy and typically developing peers. American Journal of Occupational Therapy 2009;63:96-104.
- Harding J, Harding K, Jamieson P, Mullally M, Politi C, Wong-Sing E, Law M, Petrenchik TM. Children with disabilities' perceptions of activity participation and environments: A pilot study. Canadian Journal of Occupational Therapy 2009;76:133-144.
- Beckung E, Hagberg G. Neuroimpairments, activity limitations, and participation restrictions in children with cerebral palsy. Developmental Medicine and Child Neurology 2002;44:309-316.
- Bult MK, Verschuren O, Jongmans MJ, Lindeman E, Ketelaar M. What influences participation in leisure activities of children and youth with physical disabilities? A systematic review. Research in Developmental Disabilities 2011;32:1521-1529.
- Shikako-Thomas K, Majnemer A, Law M, Lach L. Determinants of participation in leisure activities in children and youth with cerebral palsy: Systematic review. Physical and Occupational Therapy in Pediatrics 2008;28:155-169.
- 23. Palisano RJ, Chiarello LA, Orlin M, Oeffinger D, Polansky M, Maggs J, Bagley A, Gorton G. Determinants of intensity of participation in leisure and recreational activities by children with cerebral palsy. Developmental Medicine and Child Neurology 2011;53:142–149.
- Findlay LC, Garner RE, Kohen DE. Children's organized physical activity patterns from childhood into adolescence. Journal of Physical Activity and Health 2009;6:708-715.
- Statistic Sweden. Living conditions, report no 116, children's leisure time. In: Welfare S, editor. Department social welfare. Stockholm: Statistics Sweden; 2009. pp 1-54.
- Klaas SJ, Kelly EH, Gorzkowski J, Homko E, Vogel LC. Assessing patterns of participation and enjoyment in children with spinal cord injury. Developmental Medicine and Child Neurology 2010;52:468–474.
- 27. King G, McDougall J, Dewit D, Petrenchik T, Hurley P, Law M. Predictors of change over time in the activity participation of children and youth with physical disabilities. Children's Health Care 2009;38:321-351.
- King G, King S, Rosenbaum P, Kertoy M, Law M, Hurley P, Hanna S, Young N, editors. Children's assessment of participation and enjoyment & prefreences for activities of children. San Antonio, TX: Harcourt Assessment Inc; 2004.
- Guevremont A, Findlay L, Kohen D. Organized extracurricular activities of Canadian children and youth. Health Reports 2008;19:65-69.

- Kang LJ, Palisano RJ, Orlin MN, Chiarello LA, King GA, Polansky M. Determinants of social participation with friends and others who are not family members for youths with cerebral palsy. Physical Therapy 2010;90:1743-1757.
- 31. Brophy S, Cooksey R, Lyons RA, Thomas NE, Rodgers SE, Gravenor MB. Parental factors associated with walking to school and participation in organised activities at age 5: Analysis of the Millennium Cohort Study. BMC Public Health 2011;11:14.
- 32. Michelsen SI, Flachs EM, Uldall P, Eriksen EL, McManus V, Parkes J, Parkinson KN, Thyen U, Arnaud C, Beckung E, et al. Frequency of participation of 8-12-year-old children with cerebral palsy: A multi-centre cross-sectional European study. European Journal of Paediatric Neurology 2009;13:165-177.
- Beets MW, Bornstein D, Beighle A, Cardinal BJ, Morgan CF. Pedometer-measured physical activity patterns of youth: A 13-country review. American Journal of Preventive Medicine 2010;38:208-216.
- Engel-Yeger B, Jarus T, Law M. Impact of culture on children's community participation in Israel. American Journal of Occupational Therapy 2007;61:421-428.
- Mihaylov SI, Jarvis SN, Colver AF, Beresford B. Identification and description of environmental factors that influence participation of children with cerebral palsy. Developmental Medicine and Child Neurology 2004;46:299-304.
- Tisdall K. National contexts affecting the lives of disabled children in Denmark, France, Germany, Ireland, Italy, Sweden and UK (England and Northern Ireland). Newcastle: University of Newcastle upon Tyne, Robinson Library; 2006.
- Simeonsson RJ, Carlson D, Huntington GS, McMillen JS, Brent JL. Students with disabilities: A national survey of participation in school activities. Disability and Rehabilitation 2001;23:49-63.
- Imms C, Reilly S, Carlin J, Dodd KJ. Characteristics influencing participation of Australian children with cerebral palsy. Disability and Rehabilitation 2009;31:2204-2215.
- Holt L. (Dis)abling children in primary school micro-spaces: Geographies of inclusion and exclusion. Health and Place 2003;9:119-128.
- Law M, Petrenchik T, King G, Hurley P. Perceived environmental barriers to recreational, community, and school participation for children and youth with physical disabilities. Archives of Physical Medicine and Rehabilitation 2007;88:1636-1642.
- Borzecki A, Nieradko B, Gnass B, Sieklucka-Dziuba M. Leisure time activities in teenagers in urban and rural areas. Annales Universitatis Mariae Curie-Sklodowska Medicine 2002;57:375-380.
- Loucaides CA, Chedzoy SM, Bennett N. Differences in physical activity levels between urban and rural school children in Cyprus. Health Education Research 2004;19:138-147.
- 43. Imms C. Review of the Children's Assessment of Participation and Enjoyment and the Preferences for Activity of Children. Physical and Occupational Therapy in Pediatrics 2008;28:389–404.
- 44. King GA, Law M, King S, Hurley P, Hanna S, Kertoy M, Rosenbaum P. Measuring children's participation in recreation and leisure activities: Construct validation of the CAPE and PAC. Child; Care, Health and Development 2007;33:28-39.

- 45. Bult MK, Verschuren O, Gorter JW, Jongmans MJ, Piskur B, Ketelaar M. Cross-cultural validation and psychometric evaluation of the Dutch language version of the Children's Assessment of Participation and Enjoyment (CAPE) in children with and without physical disabilities. Clinical Rehabilitation 2010;24:843-853.
- Colon WI, Rodriguez C, Ito M, Reed CN. Psychometric evaluation of the Spanish version of the Children's Assessment of Participation and Enjoyment and Preferences for Activities of Children. Occupational Therapy International 2008;15:100-113.
- Ullenhag A, Almqvist L, Granlund M, Krumlinde-Sundholm L. Cultural validity of the Children's Assessment of Participation and Enjoyment/Preferences for Activities of Children (CAPE/PAC). Scandinavian Journal of Occupational Therapy 2011;16:1-11.
- Simpkins SD, Vest AE, Becnel JN. Participating in sport and music activities in adolescence: The role of activity participation and motivational beliefs during elementary school. Journal of Youth and Adolescence 2010;39:1368-1386.
- The Education Act SFS2010:800. Sweden: Ministry of Education and Research; http://www.riksdagen.se/sv/Doku ment-Lagar/Svenskforfattningssamling/Skollag-2010800_sfs-2010/ Website assessed 14062012
- The Education Act 1998: Section 5. Norway: Ministry of Education and Research; 2010; http://www.european-agency. org/country-information/norway/norwegian-files/Education_ Act Norway.pdf. Website assessed 14062012.
- 51. European Agency for Development in Special Needs Education, 2010. Special Needs Education Country Data 2010. Odense, Denmark: European Agency for Development in Special Needs Education; 2010. Available online at: http://www.european-agency.org/country-information, accessed 14062012.
- 52. Swedish Act Concerning Support and Service for Persons with Certain Functional Impairments. Swedish disability policy-service and care for people with functional impairments. http://www.socialstyrelsen.se/Lists/Artikelkatalog/attachments/8407/2009-126-188_2009126188.pdf Website assessed 14062012.
- Lane KL, Givner CC, Pierson MR. Teachers expectations of student behavior: Social skills necessary for success in elementary school classrooms. Journal of Special Education 2004;32:104-110.
- Eriksson L, Granlund M. Conceptions of participation in students with disabilities and persons in their close environment. Journal of Developmental and Physical Disabilities Applied Psychology 2004;16:229-245.
- Palisano RJ, Rosenbaum P, Bartlett D, Livingston MH. Content validity of the expanded and revised Gross Motor Function Classification System. Developmental Medicine and Child Neurology 2008;50:744-750.
- 56. Eliasson AC, Krumlinde-Sundholm L, Rosblad B, Beckung E, Arner M, Ohrvall AM, Rosenbaum P. The Manual Ability Classification System (MACS) for children with cerebral palsy: Scale development and evidence of validity and reliability. Developmental Medicine and Child Neurology 2006;48:549-554.
- 57. Hidecker MJ, Paneth N, Rosenbaum PL, Kent RD, Lillie J, Eulenberg JB, Chester K Jr., Johnson B, Michalsen L, Evatt M, et al. Developing and validating the Communication Function Classification System for individuals with cerebral palsy. Developmental Medicine and Child Neurology 2011;53:704-710.

Appendix A: Hierarchical multiple regression with Diversity outcome score as dependent variable and independent variables; age, gender, education level of mothers', rural/urban and Sweden, Norway and the Netherlands as dummy variables for children with and without disabilities.

	Children wi	th disabilities	Children with	out disabilities
Variable	Step 1	Step 2	Step 1	Step 2
Recreational		-		
Age	-0.476**	-0.478**	-0.233**	-0.228**
Gender	-0.137*	-0.148**	-0.113*	-0.113*
Educational level		-0.060	0.115	-0.115
Rural/urban		-0.011		-0.040
Sweden		-0.075	• •	
Norway		-0.073		0.115**
The Netherlands		0.166**		-0.034
R^2	0.0444	-0.166**		0.176**
	-0.241**	0.267**	0.068**	0.109**
R ² change		0.026		0.041
Sig.F change		0.066		0.000
df	251	247	425	421
F	39.776	15.005	15.448	8.578
Physical				
Age	-0.122*	-0.118*	-0.039	-0.044
Gender	0.213**	0.212**	0.257**	0.253**
Educational level		0.083		0.098
Rural/urban		-0.192**		-0.055
Sweden		0.099		-0.055
Norway		0.055		0.005
The Netherlands	•	-0.113		-0.005
R^2	0.060**		. 0.04	-0.108
R ² change	0.062**	0.121**	0.067**	0.088**
		0.080		0.020
Sig.F change		0.000		0.053
df	251	247	425	421
F	8.266	6.790	15.372	6.764
Social				
Age	-0.111	-0.105	0.026	0.015
Gender	-0.105	-0.118*	-0.176**	-0.184**
Educational level		0.021		0.077
Geographic area		-0.047		-0.035
Sweden	•	-0.110	•	-0.033
Norway		0.110		0.024
The Netherlands		<i>−0,475</i> **		
R^2	0.000			-0.214**
R ² change	0.022	0.246**	- 0.032**	0.083**
	_	0.224		0.051
Sig.F change	•	0.000		0.000
df	251	247	425	421
F	2.882	13.435	6.946	6.360
Skill-based				
Age	-0.121*	-0.099	-0.083	-0.090
Gender	-0.231**	-0.221**	-0.305**	-0.314**
Educational level		0.148*		-0.012
Geographic area		-0.044		-0.128**
Sweden	•	0.134*		-0.016
Norway		0.131		0.071
The Netherlands		-0.152*	•	
R^2	0.066**		0.101**	-0.027
R ² change	0.000	0.150**	0.101**	0.127**
		0.084		0.026
Sig.F change	051	0.000	*	0.015
df	251 ,	247	424	420
F	8.919	7.291	23.765	10.159
Self-improvement				
Age	-0.095	-0.080	-0.140**	-0.134**
Gender	-0.112	-0.113	-0.315**	-0.312**
Educational level		0.074		-0.000

(continued)

Appendix A. Continued.

		ith disabilities	Children with	out disabilities
Variable	Step 1	Step 2	Step 1	Step 2
Geographic area	,	-0.003		0.015
Sweden		0.001		
Norway				0.061
The Netherlands		-0.297**		0.045
R^2	0.021	0.120**	0.120**	0.124**
R ² change		0.099		0.004
Sig.F change		0.000		0.754
df	251	247	425	421
F	2.678	5.614	29.082	9.963

^{*}p < 0.05, ** p < 0.01 values for independent variables presented as Standardized Coefficient Beta.

Appendix B: Hierarchical multiple regression with Intensity: seldom regular and often activity outcome score as dependent variable and independent variables; age, gender, mother's educational level, rural/urban and Sweden, Norway and the Netherlands as dummy variables for children with and without disabilities.

Variable Recreational seldom	Step 1		Children without disabilities				
Recreational seldom		Step 2	Step 1	Step 2			
1 CC CUITOTH SCHOOL				•			
Age	0.476**	0.476**	0.240**	0.237**			
Gender	0.137*	0.148**	0.113*	0.114*			
Educational level		0.061		0.038			
Geographic area		-0.010		0.057			
Sweden		0.058					
Norway				0.058			
The Netherlands		0.165**		-0.149**			
R^2	0.241**	0.267**	0.071 **	0.106**			
R ² change	,	0.025		0.034			
Sig.F change		0.076		0.003			
df 2/4	251	247	· 425	421			
F	39.898	14.970	16.323	8.277			
Recreational regular		2 3.0					
Age	-0.198**	-0.184**	-0.081	-0.091			
Gender	0.029	-0.046	-0.054	-0.061			
Educational level	, 0.023	-0.094		0.063			
Geographic area	•	0.007		-0.060			
Sweden		0.023		*****			
Norway		0.023	•	-0.128*			
The Netherlands		<i>−0.318</i> **		-0.082			
R^2	0.040**	0.146**	0.010	0.032*			
R ² change	0.010	0.106	0.010	0.022			
Sig.F change		0.000	•	0.049			
df 2/4	251	247	425	421			
F	5.197	7.053	2.068	2.301			
Recreational often							
Age	-0.413**	-0.421**	-0.209**	-0.200**			
Gender	-0.135*	-0.137*	-0.088	-0.085			
Educational level	, 0.133	0.012	0.000	-0.081			
Geographic area		0.007*		-0.024			
Sweden		0.078		0.021			
Norway		0.004		0.019			
The Netherlands		0.004	• •	0.214**			
R2	0.184**	0.191**	0.052**	0.102**			
R^2 change	0.104	0.006	0.032	0.102			
Sig.F change		0.744		0.000			

(continued)

Appendix B. Continued.

		ith disabilities	Children with	out disabilities
Variable	Step 1	Step 2	Step 1	Step 2
df 2/4	251	247	425	421
F	28.238	9.711	11.709	7.986
Physical seldom		•		
Age	0.124*	0.120*	0.040	0.046
Gender	-0.211**	-0.211**	-0.265**	0.046 -0.260**
Educational level	0.211	-0.085	-0.203	-0.105*
Geographic area		0.196**		0.047
Sweden		-0.109	÷	0.047
Norway				0.025
The Netherlands		0.108	•	0.128**
R^2	0.062**	0.123**	0.071**	0.096**
R ² change		0.082		0.025
Sig.F change		0.000		0.022
df 2/4	251	247	425	421
F	8.285	6.901	16.325	7.471
Physical regular			· ·	
Age	-0.066	-0.049	0.018	A AAA
Gender	-0.042	-0.049 0.035	0.018	0.000 0.049
Educational level	-0.042	0.027	0.039	0.049
Geographic area		-0.107	•	-0.032
Sweden		0.106		-0.032
Norway		0.100		−0.110 *
The Netherlands		-0.277**		-0.110* -0.234**
R^2	0.006	0.138**	0.004	0.059**
R ² change	5.555	0.132**	7	0.055**
Sig.F change		0.000		0.000
df 2/4	251	247	425	421
F	0.795	6.587	0.809	4.365
Dl				
Physical often	0.007	0.405		
Age Gender	-0.097 <i>0.217</i> **	-0.105 0.222**	-0.059	-0.055
Educational level	0.217		0.276**	0.277**
Geographic area		0.080 0.150*		0.074
Sweden		0.048		-0.033
Norway		0.048		0.043
The Netherlands		0.080		
R^2	0.058**	0.082**	0.079**	0.003 .087**
R ² change	0.050	0.032	0.079***	0.008
Sig.F change		0.172		0.469
df 2/4	251	247	425	421
F	7.741	3.680	18.260	6.681
		21300	10.200	0.001
Social seldom	.			
Age	0.107	0.098	-0.024	-0.011
Gender	0.102	0.114*	0.177**	0.185**
Educational level		-0.026		-0.083
Geographic area		0.049		0.034
Sweden		0.088	•	
Norway The Netherlands		0 4c7++		-0.005
R^2	0.001	0.467**	0.020++	0.234**
R ² change	0.021	<i>0.240</i> ** 0.220	0.032**	0.090**
Sig.F change		0.220		0.058
df 2/4	251	247	425	0.000
ui <i>214</i> F	2.683	13.035	425 6.974	421
	2.003	13.033	0.974	6.915
Social regular				,
Age	-0.075	-0.052	-0.109*	-0.106*
Gender	-0.006	-0.002	<i>−0.128</i> **	-0.128**
Educational level		0.095		0.026

(continued)

Appendix B. Continued.

	Children wit	th disabilities	Children with	out disabilities
Variable	Step 1	Step 2	Step 1	Step 2
Geographic area		0.055		-0.040
Sweden		0.017		
Norway		•		-0.062
The Netherlands		<i>-0.310</i> **		0.075
R ²	0.006	0.116**	0.029**	0.043**
R ² change		0.111		0.015 0.164
Sig.F change	051	0.000	425	421
df 2/4 F	251 0.706	247 5.409	6.237	3.184
r	0.706	5.409	0.237	3.104
Social often				
Age	-0.067	-0.076	0.122*	0.105*
Gender	-0.121	-0.140*	-0.093	-0.103*
Educational level		-0.052		0.069
Geographic area	•	-0.108		-0.004
Sweden		-0.125*		
Norway	•			0.059
The Netherlands	ř.	-0.309 * *		-0.336**
R^2	0.019	0.139**	0.023**	0.149**
R ² change		0.120		0.126
Sig.F change		0.000	•	0.000
df 2/4	251	247	425	421
F	2.380	6.637	5.080	12.314
Skill-based seldom				
Age	0.124*	0.101	0.084	0.091*
Gender	0.232**	0.221**	0.294**	0.303**
Educational level		-0.148*		-0.007
Geographic area		0.044	,	0.113*
Sweden		-0.142*		
Norway	•		••	-0.051
The Netherlands		0.151*		0.050
R^2	0.067**	0.154**	0.094**	0.114**
R ² change		0.087		0.020
Sig.F change		0.000		0.055
df 2/4	0.0454	248	00.107	422
F	9.047**	9.020**	22.107	9.017
Skill-based regular				
Age	-0.046	-0.022	0.044	-0.048
Gender	-0.048	-0.045	<i>−0.141</i> **	-0.144**
Educational level		0.078		-0.067
Geographic area		-0.078		-0.016
Sweden		0.183**		0.070
Norway		0.164*		-0.070
The Netherlands R ²	2 224	-0.164* <i>0.101</i> **	0.022**	-0.018 0.035*
	0.004		0.022**	
R ² change		0.096		0.013 0.245
Sig.F change	051	0.000	424	
df 2/4 F	251 0.541	247 4.611	4.780	420 2.509
	0.541	4.011	4.760	2.309
Skill-based often	0.112	0.100	0.070	-0.078
Age	-0.113	-0.100 0.220**	-0.072 -0.258**	-0.078 -0.266**
Gender	-0.230**	-0.220**	-0.238	
Educational level		-0.120		0.042
Geographic area		-0.008		-0.117*
Sweden		0.062		0.024
Norway		0.000		
The Netherlands R ²	0.04**	-0.083	0.072**	-0.046 0.090**
R ² change	0.064**	0.095**	0.072^^	
л cnange		0.031		0.017

Appendix B. Continued.

	Children w	rith disabilities	Children with	out disabilities
Variable	Step 1	Step 2	Step 1	Step 2
Sig.F change	•	0.079		0.091
df 2/4	251	` 247	424	420
F	8.596	4.329	16.532	6.908
Self-improve seldom		41		
Age	0.060	0.042	0.143**	0.138**
Gender	0.102	0.106	0.314**	0.310**
Educational level		-0.062	0.514	-0.020
Geographic area		0.016		-0.020 -0.027
Sweden	•	-0.013		-0.027
Norway		-0.015		0.040
The Netherlands		0.352**		-0.040
R^2	0.014	0.352***	0.100**	-0.018
R ² change	0.014	0.131**	0.120**	0.122**
Sig.F change				0.002
df 2/4	251	0.000		0.913
GI 2/4 F	251	247		422
r	1.733	7.339	29.006	9.763
Self-improve regular		•		
Age	0.002	0.024	-0.030	-0.031
Gender	0.007	0.014	<i>−0.142</i> **	-0.140**
Educational level		0.116		0.004
Geographic area		-0.058		0.043
Sweden		0.150*		0.043
Norway				-0.061
The Netherlands		<i>−0.151</i> *	. "	-0.001 -0.017
R^2	0.000	0.082**	0.021*	0.028
R ² change	, 0.000	0.082	. 0.021	0.028
Sig.F change		0.002		
df 2/4	251	247		0.597
F	0.006	3.687	4.597	422 1.990
Self-improve often				
Age	-0.073	-0.068	-0.151**	0.14444
Gender	-0.128*	-0.138*		-0.144**
Educational level	-0.128		<i>−0.280</i> **	-0.277**
Geographic area		-0.013		0.021
Geographic area Sweden		0.025		0.003
Norway		-0.096		
5				0.088
The Netherlands R ²	0.001	-0.312**		0.032
	0.021	0.105**	0.103**	0.110**
R ² change		0.084		0.007
Sig.F change		0.000		0.505
df 2/4	251	247	425	421
F	2.705	4.831	24.283	8.637

^{*} $p \le 0.05$, ** $p \le 0.01$. Seldom or never: 1-2-times in the past 4 months; Regular: 1-3 times a month; Often: once a week-daily.

Copyright of Developmental Neurorehabilitation is the property of Taylor & Francis Ltd and its content may not be copied or emailed to multiple sites or posted to a listserv without the copyright holder's express written permission. However, users may print, download, or email articles for individual use.